

# Lecture 3

## Statistical estimation and hypothesis testing for exponential family GLM

---

# Today's topics:

- Likelihood score equation for general link
- Asymptotic distribution of the MLE estimates
- Hypothesis testing for  $\beta$
- Reading: Agresti Chapter 4.3, Faraway Chapter 8.3

# Likelihood score equation for a general link

Let  $\eta_i = g(\mu_i) = X_i^T \beta$  Then

$$\frac{\partial L_i}{\partial \beta_j} = \frac{\partial L_i}{\partial \theta_i} \frac{\partial \theta_i}{\partial \mu_i} \frac{\partial \mu_i}{\partial \eta_i} \frac{\partial \eta_i}{\partial \beta_j}$$

We have

- $\frac{\partial L_i}{\partial \theta_i} = \frac{y_i - b'(\theta_i)}{a(\phi)} = \frac{y_i - \mu_i}{a(\phi)}$
- $\frac{\partial \theta_i}{\partial \mu_i} = \frac{1}{b''(\theta_i)} = \frac{a(\phi)}{\text{Var}(y_i)}$
- $\frac{\partial \mu_i}{\partial \eta_i} = \frac{\partial \mu_i}{\partial g(\mu_i)} = \frac{1}{g'(\mu_i)}$
- $\frac{\partial \eta_i}{\partial \beta_j} = x_{ij}$

# Likelihood score equation for a general link

- The score equations can be written as

$$\frac{\partial L}{\partial \beta_j} = \sum_i \frac{(y_i - \mu_i)x_{ij}}{\text{Var}(y_i)} \frac{1}{g'(\mu_i)} = 0$$

- $\mu_i$  and  $\text{Var}(y_i)$  are both functions of  $\beta = (\beta_1, \dots, \beta_p)$
- The score equations only depend on the mean and variance of  $y_i$
- Matrix form of the score equation:

$$\dot{L}(\beta) = X^T D V^{-1} (y - \mu) = 0$$

where  $V = \text{diag}(\text{Var}(y_1), \dots, \text{Var}(y_n))$  and  $D = \text{diag}(g'(\mu_1), \dots, g'(\mu_n))^{-1}$ ,  
 $y = (y_1, \dots, y_n)$  and  $\mu = (\mu_1, \dots, \mu_n)$ .

- $L$  is not necessarily a concave function of  $\beta$

# Likelihood score equation for a general link

## Special cases

- If the link function is the canonical link, then  $D = \frac{1}{a(\phi)} V$ , thus the score equation becomes

$$\frac{1}{a(\phi)} X^T (y - \mu) = 0$$

the same as we derived earlier

- If we assume that  $g(\mu_i) = \mu_i = X_i^T \beta$ , then the estimating (score) equation becomes

$$\sum_i \frac{(y_i - X_i^T \beta) X_i}{\text{Var}(y_i)} = 0$$

which looks like weighted least square (difference: weights can depend on  $\beta$ )

# Likelihood score equation for the dispersion parameter

- The MLE estimation of  $\boldsymbol{\beta}$  for both the general and canonical link does not require knowing  $\phi$
- Statistical inference of  $\boldsymbol{\beta}$  may need an estimate of  $\phi$  (see later)
  - Example: we need to estimate  $\sigma^2$  in linear regression for calculating test statistics of the coefficients

How to estimate  $\phi$ ?

- We can also use MLE: find  $\phi$  by solving the equation:

$$\frac{\partial L}{\partial \phi} = 0$$

- $\frac{\partial L}{\partial \phi}$  also depends on  $\boldsymbol{\beta}$ : plug-in the MLE estimate  $\hat{\boldsymbol{\beta}}$
- Example: for Gaussian linear models:  $L = -\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \mathbf{X}_i^T \boldsymbol{\beta})^2 - n \log(\sqrt{2\pi}\sigma)$ 
  - $\frac{\partial L}{\partial \sigma^2} = \frac{1}{2(\sigma^2)^2} \sum_{i=1}^n (y_i - \mathbf{X}_i^T \boldsymbol{\beta})^2 - \frac{n}{2\sigma^2} \implies \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \mathbf{X}_i^T \hat{\boldsymbol{\beta}})^2$

# Statistical inference for GLM

```
## Call:
## glm(formula = y ~ weight + factor(color), family = poisson(),
##      data = Crabs)
##
## Deviance Residuals:
##      Min        1Q    Median        3Q        Max
## -2.9833  -1.9272  -0.5553   0.8646   4.8270
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  -0.04978    0.23315  -0.214   0.8309
## weight        0.54618    0.06811   8.019 1.07e-15 ***
## factor(color)2 -0.20511    0.15371  -1.334   0.1821
## factor(color)3 -0.44980    0.17574  -2.560   0.0105 *
## factor(color)4 -0.45205    0.20844  -2.169   0.0301 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
##      Null deviance: 632.79  on 172  degrees of freedom
## Residual deviance: 551.80  on 168  degrees of freedom
## AIC: 917.1
##
## Number of Fisher Scoring iterations: 6
```

- How do we get the standard error, z value and p-value of the GLM estimates?
- What does the deviance mean in this table?

# Asymptotic distribution of MLE estimation

- The MLE  $\hat{\beta}$  is consistent for the true value  $\beta_0$  when  $n \rightarrow \infty$  and  $p$  is fixed
- Asymptotic normality: when  $n$  is large

$$\hat{\beta} - \beta_0 \overset{\sim}{\sim} N(0, V_{\beta_0})$$

where  $\beta_0$  is the true value of the parameter. ( $nV_{\beta_0} = O(1)$ )

- As an applied course, we ignore the discussions of the conditions of the above consistency and CLT results, and skip the proofs.



# Calculation of $V_{\beta_0}$

- Taylor expansion (local linear approximation):

$$0 = \dot{L}(\hat{\beta}) \approx \dot{L}(\beta_0) + \ddot{L}(\beta_0)(\hat{\beta} - \beta_0)$$

- Then

$$\hat{\beta} - \beta_0 \approx - \left( \ddot{L}(\beta_0) \right)^{-1} \dot{L}(\beta_0) = - \frac{1}{\sqrt{n}} \left( \frac{\ddot{L}(\beta_0)}{n} \right)^{-1} \left( \frac{\dot{L}(\beta_0)}{\sqrt{n}} \right)$$

# Calculation of $V_{\beta_0}$

Under appropriate conditions, we have

$$\ddot{L}(\beta_0)/n = \sum_i \ddot{L}_i(\beta_0)/n \rightarrow \text{Const.} \quad (\text{law of large numbers})$$

$$\frac{\dot{L}(\beta_0)}{\sqrt{n}} = \frac{\sum_i \dot{L}_i(\beta_0)}{\sqrt{n}} \xrightarrow{d} N(0, V) \quad (\text{central limit theorem})$$

Thus we have

$$V_{\beta_0} = \left( \mathbb{E} \left( \ddot{L}(\beta_0) \right) \right)^{-1} \text{Var} \left( \dot{L}(\beta_0) \right) \left( \mathbb{E} \left( \ddot{L}(\beta_0) \right) \right)^{-1}$$

# Calculation of $V_{\beta_0}$

- The above calculation also can also be used to find the variance of  $\hat{\beta}$  from a general estimating equation  $\varphi(\hat{\beta}) = 0$  (will discuss more in later lectures)

- Property of the likelihood score equation:

Thus 
$$\text{Var} \left( \dot{L}(\beta_0) \right) = \mathbb{E} \left( \left( \left. \frac{\partial L}{\partial \beta} \right|_{\beta=\beta_0} \right)^2 \right) = -\mathbb{E} \left( \ddot{L}(\beta_0) \right)$$

- We also have 
$$V_{\beta_0} = -\mathbb{E} \left( \ddot{L}(\beta_0) \right)^{-1}$$

$$V_{\beta_0} = (X^T W X)^{-1} \text{ where } W = D^2 V^{-1}$$

- If we use a canonical link, then  $W = \frac{D}{a(\phi)} = V/a^2(\phi)$

# Asymptotic distribution of any function $h(\hat{\beta})$

- $h(\hat{\beta})$  is a consistent estimator of  $h(\beta_0)$

- We use Delta method to understand its uncertainty:

$$h(\hat{\beta}) \approx h(\beta_0) + \dot{h}(\beta_0)^T (\hat{\beta} - \beta_0)$$

$$\sqrt{n} \left( h(\hat{\beta}) - h(\beta_0) \right) \rightarrow N \left( 0, n \dot{h}(\beta_0)^T V_{\beta_0} \dot{h}(\beta_0) \right)$$

- Example: use Delta method to obtain a CI for  $\mu_i = g^{-1}(X_i^T \beta_0)$  of any individual  $i$

# Hypothesis testing

- How to test

$$H_0 : A\beta_0 = a_0 \quad V.S. \quad H_1 : A\beta_0 \neq a_0$$

- Example:  $H_0: \beta_1 = 0$  V.S.  $H_1: \beta_1 \neq 0$
- We will introduce three types of tests:
  - Wald test
  - Score test
  - Likelihood-ratio test

# Wald test

- Test statistic

$$T = (A\hat{\beta} - a_0)^T \left[ \widehat{\text{Var}}(A\hat{\beta}) \right]^{-1} (A\hat{\beta} - a_0)$$

- $\widehat{\text{Var}}(A\hat{\beta}) = AV_{\hat{\beta}}A^T$

- If  $a_0$  is a scalar, then we can rewrite the test statistic as the Wald statistic

$$z = \frac{A\hat{\beta} - a_0}{\sqrt{\widehat{\text{Var}}(A\hat{\beta})}}$$

- Under  $H_0$ , when  $n$  is large Wald statistic  $z \overset{\cdot}{\sim} N(0, 1)$

- We can also obtain a 95% CI for  $A\hat{\beta}$ :  $[A\hat{\beta} - 1.96\sqrt{\widehat{\text{Var}}(A\hat{\beta})}, A\hat{\beta} + 1.96\sqrt{\widehat{\text{Var}}(A\hat{\beta})}]$

# Wald test

- Test statistic

$$T = (A\hat{\beta} - a_0)^T \left[ \widehat{\text{Var}}(A\hat{\beta}) \right]^{-1} (A\hat{\beta} - a_0)$$

- $\widehat{\text{Var}}(A\hat{\beta}) = AV_{\hat{\beta}}A^T$

- If  $a_0$  is in general  $d$ -dimensional , then under  $H_0$ ,  $T \sim \chi_d^2$
- The Wald statistic is the “z-value” in the R GLM output for each coefficient  $\beta_j$

# A potential issue with Wald test

Let's look at an example of using Wald test for Binomial data  $y_i \sim \text{Binomial}(n_i, p_i)$  where we work on the null model:

$$\log \frac{p_i}{1 - p_i} = \log \frac{\mu_i}{n_i - \mu_i} = \beta_0$$

- We can treat the above model as using a canonical link with  $X$  being 1, then the asymptotic variance of  $\beta_0$  is

$$V_{\beta_0} = \left( \sum_i V_i \right)^{-1} = \left( \sum_i n_i p(1 - p) \right)^{-1}$$

- An estimate  $\hat{V}_{\beta_0} = V_{\hat{\beta}} = [(\sum_i n_i) \hat{p}(1 - \hat{p})]^{-1}$  where  $\hat{p}_i = \hat{p} = e^{\hat{\beta}} / (1 + e^{\hat{\beta}})$
- If we are interested in testing  $H_0 : p_i \equiv 0.5$  or equivalently  $H_0 : \beta_0 = 0$ , the Wald statistics is

$$z = \hat{\beta} \sqrt{\left( \sum_i n_i \right) \hat{p}(1 - \hat{p})}$$



# A potential issue with Wald test

- An estimate  $\widehat{V}_{\beta_0} = V_{\hat{\beta}} = [(\sum_i n_i)\hat{p}(1 - \hat{p})]^{-1}$  where  $\hat{p}_i = \hat{p} = e^{\hat{\beta}} / (1 + e^{\hat{\beta}})$
- If we are interested in testing  $H_0 : p_i \equiv 0.5$  or equivalently  $H_0 : \beta_0 = 0$ , the Wald statistics is

$$z = \hat{\beta} \sqrt{(\sum_i n_i)\hat{p}(1 - \hat{p})}$$

- Let's assume we only have one sample
  - Score equation:  $y - np = 0$ , so  $\hat{p} = y/n$
  - If  $y = 23$  and  $n = 25$ , then  $z = 3.31$
  - If  $y = 24$  and  $n = 25$ , then  $z = 3.11$ .
  - We have a smaller  $z$  value when we have stronger evidence against the null?

# A potential issue with Wald test

- On the other hand, we use the Wald test to directly test for  $H_0: p_i \equiv 0.5$
- In the example with only one sample, we can obtain the asymptotic distribution of  $\hat{p}$  directly, which results in another Wald statistic

$$z = \frac{\hat{p} - 0.5}{\sqrt{\hat{p}(1 - \hat{p})/n}}.$$

- If  $y = 23$  and  $n = 25$ , then  $z = 7.74$
  - If  $y = 24$  and  $n = 25$ , then  $z = 11.74$ .
- So the Wald statistics is not unique and depends on parameterization
  - We will discuss this more when we learn binary GLM (Chapter 5.3.3)

# Score test

- We only discuss the simple case

$$H_0 : \beta = \beta_0 \in \mathbb{R}^p \quad \text{V.S.} \quad H_1 : \beta \neq \beta_0$$

- Last time we used the property of the likelihood that:

$$\text{Var} \left( \dot{L}(\beta_0) \right) = \mathbb{E} \left( \left( \frac{\partial L}{\partial \beta} \Big|_{\beta=\beta_0} \right)^2 \right) = -\mathbb{E} \left( \ddot{L}(\beta_0) \right)$$

- The score test uses the test statistic

$$T = -\dot{L}(\beta_0)^T \left( \ddot{L}(\beta_0) \right)^{-1} \dot{L}(\beta_0)$$

and makes use of the asymptotic normal distribution of  $\dot{L}(\beta_0)$

- Under the null, we have  $T \rightarrow \mathcal{X}_p^2$  when  $n \rightarrow \infty$ .

# Likelihood ratio test

- We test for the null

$$H_0 : A\beta_0 = a_0 \quad V.S. \quad H_1 : A\beta_0 \neq a_0$$

- The likelihood ratio test statistic is

$$-2 \log \Lambda = -2 \left( L(\tilde{\beta}) - L(\hat{\beta}) \right)$$


- $\tilde{\beta}$  is the MLE of under the constraint  $A\beta = a_0$ , and  $\hat{\beta}$  is our original MLE without any constraints (under the alternative). As  $n \rightarrow \infty$ , under the null

$$-2 \log \Lambda \rightarrow \chi_d^2$$

# Comparison of the three tests

- We test for the null

$$H_0 : A\beta_0 = a_0 \quad \text{V.S.} \quad H_1 : A\beta_0 \neq a_0$$



- Three tests are asymptotically equivalent under the null
- We can also construct CI from score and likelihood ratio tests by inverting the tests