

Topics in Causal Inference

STAT41530

Jingshu Wang

Lecture 4

Topic: causal directed acyclic graph (DAG)

- Do-operator
- Confounding
 - Backdoor criterion
 - Frontdoor criterion

Do-operator

- Assume that we have a causal DAG with structural equations


$$X_j = f_j(PA_j, E_{X_j}), \quad j = 1, \dots, n$$

- Do-operator: $P(Y | do(X))$ to describe a causal effect

Definition (pearl 1995)


Given two disjoint sets of variables, X and Y , the causal effect of X on Y , denoted as $P(Y | do(X))$, is a function from X to the probability distribution on Y . For each realization x of X , $P(Y | do(X))$ gives the probability of $Y = y$ induced on deleting from the above set of structural equations all equations corresponding to variable X and substituting x for X in the remainder

$$P(X_i | do(X_j) = x) = \sum_{V/X_i} \frac{P(V)}{P(X_j | PA_j)} 1_{X_j=x}$$




$$P(X_5 | do(X_2)) ?$$

= $P(X_5 | X_2)$ in the DAG after intervention



An illustration of the do-operator



$$P(A_1, A_2, Y) = P(A_1)P(A_2|A_1)P(Y|A_1, A_2)$$

In general,

$$P(X_i = x_i | do(X_j) = x_j) = P(\underline{X_i(x_j)} = x_i)$$

- $P(Y|do(A_1, A_2)) = P(Y|A_1, A_2)$
- $P(Y|do(A_1)) = \sum_{a_2} P(A_2 = a_2|A_1)P(Y|A_1, A_2 = a_2) = P(Y|A_1)$
- $P(Y|do(A_2)) = \sum_{a_1} P(A_1 = a_1)P(Y|A_1 = a_1, A_2) \neq P(Y|A_2)$

Potential outcome

Different from conditional probability

A fundamental theorem

(Adjustment of direct causes)

Let Y be any set of variables disjoint of $\{X_i \cup PA_i\}$. The causal effect of X_i on Y is given by

$$P(Y|do(X_i)) = \sum_{pa_i} P(PA_i = pa_i)P(Y|X_i, PA_i = pa_i)$$


Where $P(PA_i = pa_i)$ and $P(Y|X_i, PA_i = pa_i)$ represent pre-interventional probabilities

Define $W = V/\{X_i, Y, PA_i\}$


(Before we change the DAG)

$$\begin{aligned}\mathbb{P}[Y | do(X_i) = x] &= \sum_W \sum_{PA_i} \sum_{X_i} \mathbb{P}[V | do(X_i) = x] \\ &= \sum_W \sum_{PA_i} \sum_{X_i} \frac{\mathbb{P}[V]}{\mathbb{P}[X_i | PA_i]} \mathbf{1}_{X_i=x} \\ &= \sum_{PA_i} \sum_{X_i} \frac{\mathbb{P}[Y, PA_i, X_i]}{\mathbb{P}[X_i | PA_i]} \mathbf{1}_{X_i=x} \\ &= \sum_{PA_i} \sum_{X_i} \frac{\mathbb{P}[Y|PA_i, X_i] \mathbb{P}[X_i | PA_i] \mathbb{P}[PA_i]}{\mathbb{P}[X_i | PA_i]} \mathbf{1}_{X_i=x} \\ &= \sum_{pa_i} \mathbb{P}[Y|PA_i = pa_i, X_i = x] \mathbb{P}[PA_i = pa_i]\end{aligned}$$

Example




DAG before intervention



DAG after intervention

$$P(X_5 | do(X_3)) = \sum_{x_1} P(X_1 = x_1) P(X_5 | X_3, X_1 = x_1)$$

Do-operator to explain Simpson's paradox



$$P(Y|do(A)) = \sum_l P(Y|L = l, A)P(L = l)$$

$$P(Y|A) = \sum_l P(Y|L = l, A)P(L = l|A)$$

$$P(Y|do(X_i)) = \sum_{pa_i} P(PA_i = pa_i)P(Y|X_i, PA_i = pa_i)$$

$$P(Y|X_i) = \sum_{pa_i} P(PA_i = pa_i|X_i)P(Y|X_i, PA_i = pa_i)$$

Structural classification of bias

Bias:




- **Systematic bias**: bias that exists even when the true distribution of observed data is known
- **Estimation bias**: bias in statistical estimation

$A \perp Y(a) \mid L$ for all a not true: introduce systematic bias

Two types of systematic bias:

- **Confounding**: treatment and outcome share a common cause
- **Selection bias**: conditioning on common effects


Confounding



If we believe that confounding is likely, how to determine L for which $A \perp Y(a) \mid L$ for all a holds?


Backdoor path: non-causal path between A and Y which has an arrow pointing to the assignment A

Examples of confounding:




Working as a firefighter \rightarrow Risk of death

Healthy working bias



Prescribe aspirin \rightarrow stroke

Confounding by indication



Linkage disequilibrium /
population stratification

The backdoor criterion

A set of covariates L satisfies the **backdoor criterion** if all backdoor paths between A and Y are blocked by conditioning on L and L contains no variables that are descendant of treatment A

Backdoor adjustment

If L satisfies the backdoor criterion, then

$$P(Y|do(A)) = \sum_l P(Y|A, L = l)P(L = l)$$

Or in other words, $A \perp Y(a) \mid L$ for all a

Proof idea of the backdoor adjustment

$$\begin{aligned} \mathbb{P}[Y \mid do(A)] &= \sum_{pa} \mathbb{P}[Y \mid A, PA = pa] \mathbb{P}[PA = pa] = \sum_{pa} \left(\sum_l \mathbb{P}[Y, L = l \mid A, PA = pa] \right) \mathbb{P}[PA = pa] \\ &= \sum_{pa} \left(\sum_l \mathbb{P}[Y \mid L = l, A, PA = pa] \mathbb{P}[L = l \mid A, PA = pa] \right) \mathbb{P}[PA = pa] \end{aligned}$$

As L satisfies the backdoor criterion:

1. L are no descents of A , PA are all direct parents of A

$$(L \perp\!\!\!\perp A \mid PA)_G \implies L \perp A \mid PA$$

2. $(L \perp\!\!\!\perp PA \mid A, L)_G \implies L \perp PA \mid A, L$

If this is not true, then there is an open path not through A and L . This introduces a backdoor path between A and Y that is not blocked by L

$$\begin{aligned} \mathbb{P}[Y \mid do(A)] &= \sum_{pa} \left(\sum_l \mathbb{P}[Y \mid L = l, A] \mathbb{P}[L = l \mid PA = pa] \right) \mathbb{P}[PA = pa] \\ &= \sum_l \mathbb{P}[Y \mid L = l, A] \sum_{pa} (\mathbb{P}[L = l \mid PA = pa] \mathbb{P}[PA = pa]) \\ &= \sum_l \mathbb{P}[Y \mid L = l, A] \mathbb{P}[L = l] \end{aligned}$$




L satisfies the backdoor criterion \implies $A \perp Y(a) \mid L$ for all a
 \impliedby (need faithfulness assumption)

Traditional way to find confounders may introduce bias

The backdoor criterion states that there is no unmeasured confounding after adjusting for L

A traditional way to find confounding covariates L

1. L is associated with A ; and
2. A is associated with Y conditional on L



In all three cases, L is not a confounder!


The frontdoor criterion

Can we identify causal effects when conditional exchangeability fails?

Definition (Pearl 1995)

A set of variables Z satisfies the front-door criterion if:

1. Z intercepts all directed paths from A to Y ; and
2. There is no unblocked backdoor path from A to Z ; and
3. All backdoor paths from Z to Y are blocked by A



Frontdoor adjustment

If Z satisfies the frontdoor criterion and if $P(A, Z) > 0$, then the causal effect of A on Y is identifiable:


$$P(Y|do(A)) = \sum_z P(Z = z|A) \sum_{a'} P(Y|A = a', Z = z)P(A = a')$$

Frontdoor adjustment


$$P(Y|do(A)) \\ = \sum_z P(Z = z|A) \sum_{a'} P(Y|A = a', Z = z)P(A = a')$$

Intuitive idea of the frontdoor adjustment:

1. $P(Z|do(A))$ is identifiable
2. $P(Y|do(Z))$ is identifiable
3. $P(Y|do(A)) = \sum_z P(Y|do(Z) = z)P(Z = z|do(A))$



Proof idea of the frontdoor adjustment



$$\begin{aligned}
 \mathbb{P}[Y \mid do(A)] &= \sum_u \mathbb{P}[Y \mid A, U = u] \mathbb{P}[U = u] = \sum_u \left(\sum_z \mathbb{P}[Y, Z = z \mid A, U = u] \right) \mathbb{P}[U = u] \\
 &= \sum_u \left(\sum_z \mathbb{P}[Y \mid Z = z, A, U = u] \mathbb{P}[Z = z \mid A, U = u] \right) \mathbb{P}[U = u]
 \end{aligned}$$

As Z satisfies the frontdoor criterion:

- $(Z \perp\!\!\!\perp U \mid A)_G \implies Z \perp U \mid A$

Otherwise, there is an unblocked backdoor path from A to Z or from Z to Y not through A

- $(A \perp\!\!\!\perp Y \mid Z, U)_G \implies A \perp Y \mid Z, U$

Z blocks paths from A and U blocks paths to A

- U blocks all backdoor paths from Z to Y

$$\begin{aligned}
 &\mathbb{P}[Y \mid do(A)] \\
 &= \sum_u \left(\sum_z \mathbb{P}[Y \mid Z = z, U = u] \mathbb{P}[Z = z \mid A] \right) \mathbb{P}[U = u] \\
 &= \sum_z \mathbb{P}[Z = z \mid A] \left(\sum_u \mathbb{P}[Y \mid Z = z, U = u] \mathbb{P}[U = u] \right) \\
 &= \sum_z \mathbb{P}[Z = z \mid A] \mathbb{P}[Y \mid do(Z)] \\
 &= \sum_z \mathbb{P}[Z = z \mid A] \left(\sum_{a'} \mathbb{P}[Y \mid Z = z, A = a'] \mathbb{P}[A = a'] \right)
 \end{aligned}$$